

PHYS-4601 Homework 16 Due 16 Feb 2012

This homework is due in the dropbox outside 2L26 by 11:59PM on the due date. If you wish to turn it in ahead of time, you may email a PDF or give a hardcopy to Dr. Frey.

1. Deformed Particle in a Box *inspired by Griffiths 6.1*

Consider a particle in a 1D infinite square well with boundaries at $x = \pm a$. If the particle is free inside the box ($-a < x < a$), the wavefunctions and energies are

$$\psi_n(x) = \left\{ \begin{array}{ll} \frac{1}{\sqrt{a}} \cos\left(\frac{n\pi}{2a}x\right) & n \text{ odd} \\ \frac{1}{\sqrt{a}} \sin\left(\frac{n\pi}{2a}x\right) & n \text{ even} \end{array} \right\}, \quad E_n^{free} = \frac{\hbar^2}{2m} \left(\frac{n\pi}{2a}\right)^2. \quad (1)$$

In the following, use first-order perturbation theory in the case that the particle is not free inside the box.

(a) For potential (inside the box)

$$V(x) = \left\{ \begin{array}{ll} V_0 & -a/2 < x < a/2 \\ 0 & |x| \geq a/2 \end{array} \right., \quad (2)$$

find the approximate ground state energy. Write this energy as E_1^{free} times a function of $\epsilon = ma^2V_0/\hbar^2$.

(b) For potential (inside the box)

$$V(x) = \alpha\delta(x) \quad (3)$$

find the approximate energies for all the states. Show that the first-order corrections to the energy are small only when $ma\alpha/\hbar^2 \ll 1$.

2. Changing Spring Constant *Griffiths 6.2 extended*

The harmonic oscillator potential is $V = (1/2)kx^2 \equiv (1/2)m\omega^2x^2$, and we've studied the states, energy levels, and ladder operators quite a bit before. Now consider a second harmonic oscillator with spring constant $k' = k(1 + \epsilon)$. *Instructions:* In this problem, do not use the wavefunction or carry out any integrals. Use the ladder operators; they will save you a lot of work.

- Suppose that $\epsilon \ll 1$. Write the energy eigenvalues of the second oscillator in terms of k and ϵ first as an exact expression and then as a power series in ϵ to second order.
- Write the potential V' for the second harmonic oscillator approximately as the first harmonic oscillator potential plus a perturbation which is first order in ϵ . Using that expression, find the difference in energies between the two harmonic oscillators at first order in ϵ using perturbation theory and compare that to your answer from part (a).
- Finally, consider the ground states of the two operators, which we can call $|0\rangle$ and $|0'\rangle$. To first order in epsilon, find the difference in these two states (which is itself an unnormalized ket).

3. Stark Effect *based on Griffiths 6.36*

The presence of an external electric field $E_0\hat{z}$ shifts the energy levels of a hydrogen atom, which is called the Stark effect. Consider the hydrogen atom to be described by the Coulomb potential; the external electric field introduces a perturbation

$$H_1 = eE_0z = eE_0r \cos \theta. \quad (4)$$

We have already seen on homework that the expectation value of this Hamiltonian in the ground state $n = 1$ vanishes, so there is no shift in the ground state energy. In this problem, we consider the degenerate perturbation theory of the $n = 2$ states. *As spin does not enter, do not consider it in this problem.*

- (a) The four states $|2, 0, 0\rangle$, $|2, 1, 0\rangle$, and $|2, 1, \pm 1\rangle$ are degenerate at 0th order. Label these states sequentially as $i = 1, 2, 3, 4$. Show that the matrix elements $W_{ij} = \langle i|H_1|j\rangle$ form the matrix

$$W = -3aeE_0 \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (5)$$

where empty elements are zero and a is the Bohr radius. *Hint:* You may use the results for the Jan 31 in-class test, problem 8, as given in the solution set. Also, note that L_z commutes with H_1 , so only states with the same quantum number m can have nonzero matrix elements.

- (b) Diagonalize this matrix to show that $|\pm\rangle = (1/\sqrt{2})(|2, 0, 0\rangle \pm |2, 1, 0\rangle)$ are eigenstates of W . Find the first order shift in energies of $|\pm\rangle$. *Hint:* You may still want that exam solution. Note that the corrected eigenstates may still have contributions from other values of the principal quantum numbers n , but that doesn't quite matter.
- (c) Finally, show that the states $|\pm\rangle$ have a nonzero dipole moment $p_z = -e\langle z\rangle$ and calculate it. You should not need to do any more calculations; just use your answer from part (b).