

PHYS-4601 Homework 9 Due 14 Nov 2013

This homework is due in the dropbox outside 2L26 by 11:59PM on the due date. If you wish to turn it in ahead of time, you may email a PDF or give a hardcopy to Dr. Frey.

1. Raising and Lowering

- (a) *More or less Griffiths 4.18* Using the relation for $L_{\pm}L_{\mp}$ given in class and the text, show that

$$L_{\pm}|\ell, m\rangle = \hbar\sqrt{(\ell \mp m)(\ell \pm m + 1)}|\ell, m \pm 1\rangle . \quad (1)$$

- (b) In a vector/matrix representation of the $\ell = 1$ states where

$$|1, 1\rangle = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} , \quad |1, 0\rangle = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} , \quad |1, -1\rangle = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} , \quad (2)$$

use (1) to find matrix representations of L_{\pm} and then L_x and L_y .

- (c) *Griffiths 4.22(b)* Use $L_{+} \cdot Y_{\ell}^{\ell} = 0$ and $L_z \cdot Y_{\ell}^{\ell} = \ell\hbar Y_{\ell}^{\ell}$ to determine $Y_{\ell}^{\ell}(\theta, \phi)$ up to overall normalization.

2. Probabilities and Expectations based on problems by Ohanian

Some particle has a wavefunction

$$\psi(\vec{x}) = \frac{1}{4}\sqrt{\frac{5}{\pi}}\sin^2\theta \left[1 + \sqrt{14}\cos\theta\right] \cos(2\phi)R(r) , \quad (3)$$

where R is normalized ($\int_0^{\infty} dr r^2 |R|^2 = 1$).

- (a) If you measure the total orbital angular momentum \vec{L}^2 , what are the possible values you would find and the probabilities you would measure each value?
- (b) Find $\langle L_x \rangle$, $\langle L_y \rangle$, and $\langle L_z \rangle$ in this state.
- (c) Find the uncertainty of L_z .

3. Landau Levels from Griffiths 4.60

This problem considers the motion of electrons which are essentially confined to a 2D surface in the presence of an orthogonal magnetic field. This is the system used to describe the quantum Hall effect. Since this is a 2D problem, we won't include p_z (if you like, you can imagine that we consider only eigenstates of p_z with zero eigenvalue).

- (a) Show that a magnetic field $\vec{B} = B_0\hat{k}$ can be described by vector potential $\vec{A} = (B_0/2)(x\hat{j} - y\hat{i})$. ($\hat{i}, \hat{j}, \hat{k}$ are unit vectors along x, y, z respectively.)
- (b) We saw on the last assignment that the Hamiltonian is

$$H = \frac{1}{2m} (\vec{p} - q\vec{A})^2 . \quad (4)$$

In this case, show that we can write

$$H = \frac{1}{2m} (p_x^2 + p_y^2) + \frac{1}{2}m\omega^2 (x^2 + y^2) - \omega L_z , \quad (5)$$

where $\omega = qB_0/2m$. Argue that $[H, L_z] = 0$ (you may use results from earlier problems).

- (c) Except for the L_z term at the end, this looks like a harmonic oscillator in the x and y directions. Write H and L_z in terms of the raising and lowering operators $a_x^\dagger, a_y^\dagger, a_x, a_y$ of those two harmonic oscillators. Evaluate $L_z|n_x, n_y\rangle$; is it diagonal?
- (d) Apparently we have not yet found how to diagonalize H and L_z simultaneously. Now define lower operators

$$A = \frac{1}{\sqrt{2}}(a_y + ia_x), \quad \bar{A} = \frac{1}{\sqrt{2}}(a_y - ia_x) \quad (6)$$

and their adjoints, the raising operators. First, show that A, A^\dagger and \bar{A}, \bar{A}^\dagger satisfy the usual commutation relations for raising and lowering operators. Then, find H and L_z in terms of $A, \bar{A}, A^\dagger, \bar{A}^\dagger$. From those expressions, argue that the energy eigenvalues are $E_n = \hbar\omega_B(n + 1/2)$, where $\omega_B = 2\omega = qB_0/m$ is the *cyclotron frequency*, and that the energy eigenstates are infinitely degenerate. These energy levels are called *Landau levels*; in practice, the finite size of the metal where the electrons live reduces the degeneracy to a finite amount.