

Quantum Mechanics I PHYS-3301 Final Exam

Dr. Andrew Frey

12 April 2016, 1:30-4:30PM, 3M67

Instructions:

- Do not turn over until instructed. You will have 3 hours to complete this exam.
- No electronic devices or hardcopy notes are allowed.
- INSTRUCTIONS ABOUT THE QUESTIONS WILL GO HERE.
- **Only the lined pages of your exam book will be graded. Use the blank pages for scratch work only.**

Useful Concepts & Formulae:

- Physical Constants

- Speed of light $c = 3 \times 10^8$ m/s = 1 lightsecond/second
- Planck constant $h = 2\pi\hbar = 7 \times 10^{-34}$ Js
- Boltzmann constant $k_B = 10^{-23}$ J/K

- Quantum Mechanics

- Hydrogen wavefunctions are described by n, ℓ, m and will be given if needed
Energy levels are $E = -13.6$ eV/ n^2
- Spherical harmonics will be given if needed
- Momentum space wavefunction $\tilde{\psi}(p)$ with $p \cdot \tilde{\psi} = p\tilde{\psi}$, $x \cdot \tilde{\psi} = i\hbar d\tilde{\psi}/dp$

$$\tilde{\psi}(p) = \int \frac{dx}{\sqrt{2\pi\hbar}} e^{-ipx/\hbar} \psi(x), \quad \psi(x) = \int \frac{dp}{\sqrt{2\pi\hbar}} e^{ipx/\hbar} \tilde{\psi}(p)$$

- Maxwell-Boltzmann, Fermi-Dirac, Bose-Einstein distributions

$$n_{MB}(E) = e^{-(E-\mu)/k_B T}, \quad n_{FD}(E) = \frac{1}{e^{(E-\mu)/k_B T} + 1}, \quad n_{BE}(E) = \frac{1}{e^{(E-\mu)/k_B T} - 1}$$

- Free particles: $E_k = \hbar^2 \vec{k}^2 / 2m$ with degeneracy $d_k = (sV/2\pi^2) k^2 dk$ where s = number of polarizations

- Galilean Relativity/Newtonian Mechanics

- Galilean boost $\vec{x}' = \vec{x} - \vec{v}t$, $\vec{u}' = \vec{u} - \vec{v}$, $\vec{p}' = \vec{p} - m\vec{v}$, $k' = k - \vec{p} \cdot \vec{v} + (1/2)mv^2$
- Kinetic energy for many particles $K = K_{int} + (1/2)MV^2$

$$M = \sum m_i, \quad \vec{V} = \frac{1}{M} \sum m_i \vec{u}_i$$

- For two particles $K_{int} = (1/2)\mu u^2$ for relative velocity \vec{u} and reduced mass $\mu = m_1 m_2 / M$

- 4-vectors and Lorentz transformations

- The position 4-vector is x^μ with $x^0 = ct$.

- The metric $\eta_{\mu\nu}$ can be written as a diagonal matrix with diagonal elements $-1, 1, 1, 1$, and the invariant interval is $ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu = -c^2 dt^2 + d\vec{x}^2 = -c^2 d\tau^2$.
- The Lorentz boost transformations (in standard configuration) are

$$t' = \gamma(t - vx/c^2), \quad x' = \gamma(x - vt), \quad y' = y, \quad z' = z, \quad \gamma = 1/\sqrt{1 - v^2/c^2}.$$

They can be written as $x^{\mu'} = \Lambda^{\mu'}_{\nu} x^\nu$

- Lowered indices $a_\mu = \eta_{\mu\nu} a^\nu$ (both in frame S)
 - Relativistic dot product $a \cdot b = \eta_{\mu\nu} a^\mu b^\nu = a_\mu b^\mu = -a^0 b^0 + \vec{a} \cdot \vec{b}$
 - Tensor transformation $T_{\mu' \dots \nu' \dots} = (\Lambda^{\alpha}_{\mu'} \dots) (\Lambda^{\nu'}_{\beta} \dots) T_{\alpha \dots \beta \dots}$
- Velocities and Momenta
 - For normal velocity $\vec{u} = d\vec{x}/dt$, the Lorentz transformation in standard configuration is

$$u'_x = \frac{u_x - v}{1 - vu_x/c^2}, \quad u'_{y,z} = \frac{u_{y,z}}{\gamma(v)(1 - vu_x/c^2)}.$$

- 4-velocity: $U^\mu = dx^\mu/d\tau$, where τ is proper time along the worldline. $U^0 = \gamma c$, $\vec{U} = \gamma d\vec{x}/dt$, $d\vec{x}/dt = c(\vec{U}/U^0)$.
- 4-momentum is $p^\mu = mU^\mu$. Energy $E = cp^0$ and momentum is the spatial part \vec{p} .
- $U_\mu U^\mu = -c^2$ and $p_\mu p^\mu = -m^2 c^2$ for a normal massive particle.

- The Doppler effect, in terms of the rest frame of the receiver, is

$$\frac{\omega_R}{\omega_E} = \frac{\sqrt{1 - u_E^2/c^2}}{1 - \hat{k} \cdot \vec{u}_E/c},$$

where \hat{k} is the direction of travel of light and \vec{u}_E is the velocity of emitter relative to receiver.

- Math

- Exponential integrals (for $n = 0, 1, \dots$ and $a > 0$)

$$\int_0^\infty dx x^n e^{-x/a} = n! a^{n+1}$$

- Gaussian integrals

$$\int_{-\infty}^\infty dx e^{-ax^2} = \sqrt{\frac{\pi}{a}}, \quad \int_{-\infty}^\infty dx x^2 e^{-ax^2} = \frac{1}{2a} \sqrt{\frac{\pi}{a}}$$

- Hyperbolic trig functions: $d \cosh \theta / d\theta = \sinh \theta$, $d \sinh \theta / d\theta = \cosh \theta$

$$\cosh^2 \theta - \sinh^2 \theta = 1, \quad \cosh^2 \theta + \sinh^2 \theta = \cosh(2\theta), \quad 2 \sinh \theta \cosh \theta = \sinh(2\theta)$$

- Binomial expansion $(1 + x)^n \approx 1 + nx$ for $x \ll 1$