

PHYS-4601 Homework 2 Due 24 Sept 2015

This homework is due in the dropbox outside 2L26 by 11:59PM on the due date. You may alternately email a PDF (typed or black-and-white scanned) or give a hardcopy to Dr. Frey.

1. Diagonalization *Based on Griffiths A.26*

Consider a three-dimensional Hilbert space with orthonormal basis $|e\rangle_i$, $i = 1, 2, 3$. The operator A takes the matrix representation

$$A = \sum_{i,j} |e_i\rangle\langle e_i| A |e_j\rangle\langle e_j| \simeq \begin{bmatrix} 2 & i & 1 \\ -i & 2 & i \\ 1 & -i & 2 \end{bmatrix}. \quad (1)$$

You should be able to check yourself that A is Hermitian.

- Find the eigenvalues a_i and corresponding eigenstates $|a_i\rangle$ ($A|a_i\rangle = a_i|a_i\rangle$) written in terms of their components $\langle e_j|a_i\rangle$. Choose the eigenstates to form an orthonormal eigenbasis; that is, choose any ambiguities such that $\langle a_i|a_j\rangle = \delta_{ij}$.
- As we will state in class, A can be written in the form

$$A = \sum_i a_i |a_i\rangle\langle a_i|, \quad (2)$$

where a_i are the eigenvalues and $|a_i\rangle$ are the eigenvectors of A . Verify that formula (2) gives the same operator as (1) when you plug in your answer to part (a) for the eigenvalues and eigenvectors.

2. Something We'll Call "Momentum"

Define a "momentum" operator p that acts as $p \simeq -i\hbar d/dx$ in the position basis (precisely, $\langle x|p|\psi\rangle = -i\hbar d\psi/dx$ for any state $|\psi\rangle$ with wavefunction $\langle x|\psi\rangle = \psi(x)$). In this problem, we will explore some properties of this operator; in the future, we will see why it makes sense to call it momentum. For simplicity, we work in one dimension.

- Let the state $|p\rangle$ be an eigenstate of the momentum operator with eigenvalue p (ie, $p \cdot |p\rangle = p|p\rangle$). Show that $|p\rangle$ has wavefunction

$$\langle x|p\rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{ipx/\hbar} \quad (3)$$

(you may assume the normalization constant is given).

- Show that $\langle p'|p\rangle = \delta(p - p')$. *Hint:* You may find the formula

$$\delta(z) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{ikz} \quad (4)$$

helpful.

- Show that the wavefunction $\psi(x) = \langle x|\psi\rangle$ and "momentum-space wavefunction" $\tilde{\psi}(p) = \langle p|\psi\rangle$ for any state ψ are Fourier transforms, as defined in Griffiths equation [2.102] (up to factors of \hbar). To work this out precisely, it will be helpful for you to rescale x and p to remove explicit powers of \hbar .

- (d) Show first that $\langle p|p|\psi\rangle = p\tilde{\psi}(p)$ for any state $|\psi\rangle$. Then use your previous result to demonstrate that

$$\langle x|p \cdot \psi\rangle = -i\hbar \frac{d\psi}{dx}(x) . \quad (5)$$

What this means is that defining $p \simeq -i\hbar d/dx$ is equivalent to defining the state $|p\rangle$ by (3) — you can derive one statement from the other.

3. Permutation Operator

Consider an N -dimensional Hilbert space with orthonormal basis $\{|1\rangle, |2\rangle, \dots, |N\rangle\}$ and define the permutation operator S such that $S|n\rangle = |n+1\rangle$ for $1 \leq n < N$ and $S|N\rangle = |1\rangle$.

- (a) Show that the state

$$|\lambda\rangle = \sum_{n=1}^N \lambda^{-n+1} |n\rangle \quad (6)$$

is an eigenstate of S with eigenvalue λ as long as λ takes one of N allowed values. Find those allowed values.

- (b) Is S ever a Hermitian operator? If so, what are the values of N such that S is Hermitian?
(c) In the orthonormal basis described, write S in matrix form for the cases of $N = 2$ and $N = 3$.