

PHYS-4601 Homework 9 Due 19 Nov 2015

This homework is due in the dropbox outside 2L26 by 11:59PM on the due date. You may alternately email a PDF (typed or black-and-white scanned) or give a hardcopy to Dr. Frey.

1. Isotropic Harmonic Oscillator *from Griffiths 4.38,39*

Consider a harmonic oscillator where the restoring force is independent of the direction. In this case, the potential is

$$V(r) = \frac{1}{2}m\omega^2 r^2 . \quad (1)$$

- Show that the energy eigenvalues are $E_n = \hbar\omega(n + 3/2)$, where n is any non-negative integer. It's easiest to do this using separation of variables in Cartesian coordinates.
- Find the degeneracy of states with energy E_n .
- Now consider the Schrödinger equation in spherical coordinates and restrict to the case $\ell = 0$ for the radial equation. We know that the ground state has $\psi \propto \exp[-\rho^2/2]$ from separation of variables in Cartesian coordinates, where $\rho = \sqrt{m\omega/\hbar} r$. Therefore, we define $u = v(\rho) \exp[-\rho^2/2]$, where $v(\rho) = \rho + \dots$ is a polynomial. Argue that the (unnormalized) wavefunctions $u = H_n(\rho) \exp[-\rho^2/2]$ solve the radial equation for any odd n . Find the associated energies for $n = 1, 3$. Did you find the energies you expected from part (a)? Explain why or why not.

2. A Finite Spherical Box *extended from Griffiths 4.9*

Consider a particle of mass m in the spherically symmetric potential

$$V(r) = \begin{cases} -V_0 & (r < a) \\ 0 & (r \geq a) \end{cases} . \quad (2)$$

In 1D quantum mechanics, any potential that goes to zero at infinity and is negative anywhere has at least one bound state. We will see that is not true in 3D.

- Assume $\ell = 0$ and energy $E < 0$. Find a transcendental equation that determines E . What is the condition on V_0 that allows a bound state?
- Use Maple to solve the transcendental equation of part (a) and plot the ground state energies as a function of V_0 in the range $\pi^2\hbar^2/8ma^2 < V_0 < \pi^2\hbar^2/2ma^2$. Attach a printout of your code and the plot. *Hint:* One way to proceed is to solve for the energy with a particular value of V_0 using `fsolve` and step through values of V_0 using `seq`, then `listplot`.
- Use the numerical method of assignment 7 to verify the bound state energy for $V_0 = \pi^2\hbar^2/4ma^2$. Plot the stationary state wavefunction in this case; note that the boundary conditions here should be similar to the 1st excited state from that assignment. Attach a copy of your Maple code (only for your final energy and wavefunction, please).

3. Electromagnetic Gauge Transformations *Griffiths 4.61*

Now that we're in 3D, we could imagine having an electromagnetic field. For a particle of charge q in potential Φ and vector potential \vec{A} , the Hamiltonian is

$$H = \frac{1}{2m} (\vec{p} - q\vec{A})^2 + q\Phi . \quad (3)$$

The electric and magnetic field are

$$\vec{E} = -\vec{\nabla}\Phi - \frac{\partial\vec{A}}{\partial t}, \quad \vec{B} = \vec{\nabla} \times \vec{A}. \quad (4)$$

For more details, see Griffiths problem 4.59.

- (a) Show that the electromagnetic fields are invariant under *gauge transformations*. That is, show that the potentials

$$\Phi' = \Phi - \frac{\partial\Lambda}{\partial t}, \quad \vec{A}' = \vec{A} + \vec{\nabla}\Lambda \quad (5)$$

give the same \vec{E} and \vec{B} fields as Φ and \vec{A} , where Λ is any function of \vec{x} and t .

- (b) Since the Hamiltonian involves the potentials, it looks like we can't just make a gauge transformation in the quantum theory. However, assuming that a wavefunction $\Psi(\vec{x}, t)$ solves the time-dependent Schrödinger equation for potentials Φ and \vec{A} , show that

$$\Psi' = e^{iq\Lambda/\hbar}\Psi \quad (6)$$

solves the time-dependent Schrödinger equation for the potentials Φ' and \vec{A}' given in (5).

This gauge invariance is a critical feature of the quantum mechanical theory of electromagnetism with profound consequences. We may explore aspects of it again in assignments.