

PHYS-4601 Homework 18 Due 16 Mar 2017

This homework is due in the dropbox outside 2L26 by 10:59PM on the due date. You may alternately email a PDF (typed or black-and-white scanned) or give a hardcopy to Dr. Frey.

1. Anharmonic Oscillator

Consider a particle moving in the potential

$$V(x) = \frac{1}{2}m\omega^2x^2 + gx^3, \quad (1)$$

where g is considered to be small, so this potential can be treated as a perturbation of a harmonic oscillator. Find the correction to the energy of the harmonic oscillator ground state $|0\rangle$ at both first and second order in g .

2. Fermi's Golden Rule

Consider a sinusoidal perturbation Hamiltonian $H_1 = Ve^{-i\omega t} + V^\dagger e^{i\omega t}$. In the class notes, we found the probability for a transition from state $|1\rangle$ to $|2\rangle$ as a function of time and frequency ω . In the following, define $\hbar\omega_0 = E_2 - E_1$, the difference of the energy eigenvalues of the unperturbed Hamiltonian H_0 . We will investigate the transition probability near $\omega = \omega_0$ at large t (at least as long as the probability stays small).

- At a fixed (and large) time, the probability is peaked at $\omega = \omega_0$. Using L'Hospital's rule or just a power series expansion, find the peak transition probability as a function of time.
- Find the values of ω where the probability first vanishes on either side of $\omega = \omega_0$. The difference in these two values tells us the width of the peak.
- For large enough times, approximate the transition probability as a rectangle with the peak value from part (a) and width given by half the difference in part (b). Integrate this approximate probability function and argue that

$$P \rightarrow \frac{2\pi|V_{21}|^2}{\hbar^2} t \delta(\omega_0 - \omega) \quad (2)$$

as $t \rightarrow \infty$.

This problem shows two things: first, transitions occur only to states at energies related by the perturbation frequency and, second, that there is a constant transition rate (probability per unit time) to the appropriate states. The relationship (2) is known as *Fermi's Golden Rule*. (There is of course a more rigorous derivation possible.)

3. Magnetic Resonance Spin Flips

Consider a spin-1/2 particle (for example, a proton) with gyromagnetic ratio γ in the presence of a magnetic field

$$\vec{B} = B_0\hat{z} + B_1 \cos(\omega t)\hat{x} - B_1 \sin(\omega t)\hat{y} \quad (3)$$

at its fixed position. This is a magnetic field with a fixed z component and another component rotating in the x, y plane.

- Write the Hamiltonian either as a matrix or in terms of spin operators and show that it takes the form $H = H_0 + Ve^{-i\omega t} + V^\dagger e^{i\omega t}$.

- (b) Assume that the rotating field B_1 is much smaller than B_0 . If the spin is initially spin up at $t = 0$, find the transition probability to spin down at a later time t using perturbation theory. *Hint:* Consider the states in the Hamiltonian H_0 and their energy differences first.
- (c) It is also possible to find this transition probability exactly. With the initial conditions given in part (b), the solution of the time-dependent Schrödinger equation is

$$\begin{aligned}\langle +|\Psi(t)\rangle &= e^{i\omega t/2} \left[\cos(\alpha t/2) - i \frac{(\omega - \gamma B_0)}{\alpha} \sin(\alpha t/2) \right] \\ \langle -|\Psi(t)\rangle &= i e^{-i\omega t/2} \frac{\gamma B_1}{\alpha} \sin(\alpha t/2)\end{aligned}\tag{4}$$

with $\alpha = \sqrt{\gamma^2 B_1^2 + (\omega - \gamma B_0)^2}$. Now use Maple to verify that (4) solves the Schrödinger equation. Input the Schrödinger equation and initial conditions as a list of equations and then the solution above as another list. Then use the `odetest` function in Maple to check that (4) solves the time-dependent Schrödinger equation. Include a copy of your Maple code.

- (d) Use (4) to find the transition probability from spin up ($|+\rangle$) to spin down ($|-\rangle$). Find the conditions that this probability is one. Finally, show that it reduces to the perturbation theory result when $\gamma B_1 \ll \omega - \gamma B_0$.

4. Variational Principle for the Linear Well

Consider a particle moving in 1D in a potential $V(x) = \alpha|x|$. Find the best possible upper bound on the ground state energy using a gaussian trial wavefunction.