

PHYS-4602 Homework 8 Due 13 March 2024

This homework is due to <https://uwcloud.uwinnipeg.ca/s/FFJiJMnt9Czgo72> by 10:59PM on the due date. Your file(s) must be in PDF format; they may be black-and-white scans or photographs of hardcopies (all converted to PDF), PDF prepared by LaTeX, or PDF prepared with a word processor *using an equation editor*.

1. Relativistic Harmonic Oscillator

Recall that the relativistic energy is $\sqrt{(\vec{p}c)^2 + (mc^2)^2} \approx mc^2 + \vec{p}^2/2m - \vec{p}^4/8m^3c^2$ (plus higher-order corrections), so a 1D harmonic oscillator with the first relativistic correction has the Hamiltonian

$$H = \frac{p^2}{2m} - \frac{p^4}{8m^3c^2} + \frac{1}{2}m\omega^2x^2. \quad (1)$$

- Find the ground state energy of this oscillator using first-order perturbation theory. What condition must the frequency satisfy for the relativistic correction to be small?
- Find the ground state eigenstate of this oscillator in terms of the unperturbed oscillator eigenstates using first-order perturbation theory.

2. Stark Effect based on *G&S 7.45*

The presence of an external electric field $E_0\hat{z}$ shifts the energy levels of a hydrogen atom, which is called the Stark effect. Consider the hydrogen atom to be described by the Coulomb potential; the external electric field introduces a perturbation

$$H_1 = eE_0z = eE_0r \cos \theta. \quad (2)$$

We have already seen on homework that the expectation value of this Hamiltonian in the ground state $n = 1$ vanishes, so there is no shift in the ground state energy. In this problem, we consider the degenerate perturbation theory of the $n = 2$ states. *As spin does not enter, do not consider it in this problem.*

- The four states $|2, 0, 0\rangle$, $|2, 1, 0\rangle$, and $|2, 1, \pm 1\rangle$ are degenerate at 0th order. Label these states sequentially as $i = 1, 2, 3, 4$. Show that the matrix elements $W_{ij} = \langle i|H_1|j\rangle$ form the matrix

$$W = -3aeE_0 \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (3)$$

where empty elements are zero and a is the Bohr radius. *Hint:* Note that L_z commutes with H_1 , so only states with the same quantum number m can have nonzero matrix elements; this will save you quite a bit of work. Then use the angular wavefunctions to see that all the diagonal elements of W must vanish. Finally, use the explicit wavefunctions to evaluate the remaining matrix elements of W (there should only be one independent one left).

- Diagonalize this matrix to show that $|\pm\rangle = (1/\sqrt{2})(|2, 0, 0\rangle \pm |2, 1, 0\rangle)$ are eigenstates of W . Find the first order shift in energies of $|\pm\rangle$. *Hint:* Note that the corrected eigenstates may still have contributions from other values of the principal quantum numbers n , but that doesn't quite matter.

- (c) Finally, show that the states $|\pm\rangle$ have a nonzero dipole moment $p_z = -e\langle z\rangle$ and calculate it. You should not need to do any more calculations; just use your answer from part (b).

3. Localized Magnetic Field

Two electrons are localized at well-separated lattice sites, so they can be treated as distinguishable particles. The two electrons interact with each other, and only the first electron experiences a magnetic field. The Hamiltonian is $H = A\vec{S}_1 \cdot \vec{S}_2 + BS_{1,z}$, where \vec{S}_j is the spin of electron j and A, B are constants. (The B term represents the magnetic field on the first electron.)

- (a) Assume that there is no magnetic field, so $B = 0$. Find the energy eigenstates and eigenvalues. Which eigenstate is the ground state assuming $A > 0$?
- (b) Assume that the magnetic field on electron 1 is small ($B \ll \hbar A$) and find the ground state and ground state energy to first order in B . *Hint:* make sure you can write your eigenstates from the previous part in terms of the individual spins.